

Correlation between life style and gingival health in adult patients in Erbil city-Iraq

Shahida Rassul Hussein

Department of Dental Assistant, Erbil Technical Medical Institute, Erbil Polytechnic University, Erbil, Iraq, shahida.husain@epu.edu.iq

Follow this and additional works at: <https://polytechnic-journal.epu.edu.iq/home>

How to Cite This Article

Hussein, Shahida Rassul (2023) "Correlation between life style and gingival health in adult patients in Erbil city-Iraq,"

Polytechnic Journal: Vol. 13: Iss. 1, Article 9.

DOI: <https://doi.org/10.59341/2707-7799.1733>

This Original Article is brought to you for free and open access by Polytechnic Journal. It has been accepted for inclusion in Polytechnic Journal by an authorized editor of Polytechnic Journal. For more information, please contact polytechnic.j@epu.edu.iq.

RESEARCH ARTICLE

correlation between life style and gingival health in adult patients in Erbilcity-Iraq

Shahida Rassul Hussein

Erbil poly technique university, medical technical institute, dentalassistant department

***Corresponding author:**

Hussein SR: Assist. Prof.:

Erbil poly technique university, medical technical institute, dental assistant department

E-mail: shahida.husain

@epu.edu.iq

Received: 27 September 2021

Accepted: 14 August 2022

Published: 20 September 2023

DOI: <https://doi.org/10.59341/2707-7799.1733>

ABSTRACT

Background and objective: Several factors may affect gingival and oral health these factors may include smoking, diet, physical activities and oral hygiene The purpose of present study was to determine the correlation between gingival health and lifestyle among adult individuals in the city of Erbil.

Method: The correlation between lifestyle and gum health were determined in 378 patients, their ages were ≥ 18 and both genders who attended medical centers in Erbil. The individuals interviewed by researcher using a researcher-designed questionnaire. The medical examinations were included: plaque index (Silness J & Loe, 1964) gingival index (Loe H & Silness, 1963) examinations of teeth were selected by (Ramfjord SP, 1959) were used.

Results: That has been proven; the highest mean value for plaque (PLI)(0.867), gingival scores(GI) (0.942) increased by increasing consumption of vegetables per day with statistically significant differences (p value = 0.000). Mean plaque (0.573) gingival score (0.673) increased with the increasing fruit consumption per day with statistically significant differences for both scores (P value = 0.000). Mean plaque (1.155) and gingival (1.010) The indices were higher in subjects ingested eggs and butter once a day, with statistically highly significant differences (p-value = 0.000). Also mean plaque (1.119), gingival scores (1.14) observed more among meat consumption once a week with a statistically highly significant difference (p-value = 0.000). mean plaque value (0.755) observed higher in the non-smoking groups without statistically significant differences (p- value = 0.149) and gingival scores (0.860), among the heavy smokers groups (p value = 0.041). Mean plaque (0.640) and gingiva (0.639) index were low in subjects whom perform daily exercise (p- value = 0.069) (p- value = 0.049) for plaque score and gingival score. Mean plaque (0.528) and gingival (0.598) indices were low in people whom performing their daily tooth brushing with statistical significant difference (P value = 0.011) plaque score and statistical highly significant difference (P value = 0.001) for gingival score.

Conclusions: There was a positive correlation between lifestyle and gingival health

Key words: Gingival health, smoking, lifestyle, exercise and diet

INTRODUCTION

Background

Periodontitis is characterized by microbial-associated, host mediated inflammation that results in loss of periodontal attachment. (Maurizio S T et al , 2018) . Periodontal and disease of the gums, with their inflammatory process, steadily spread, and increase by way of the presence of dental plaque and tartar. (Corbet E, 2007).Marginal gingival Inflammation is not unusual circumstance the volume and severity can be variable. Gingival inflammation that produce by plaque may be changed by means of systemic and local affects. gingivitis can be reversed if oral hygiene are delivered. (Martha E. Nunn,2000,2003) Numerous research has been determined periodontal diseases and the effect of plaque as etiological factor. (Albander JM,2000,2002), (Loe H,2000). The most broadly danger thing is dental plaque that accumulate on the enamel due to the lack of oral

hygiene. (LoeH,2000), (Pihlstrom BL,2005).

Lifestyles commonly manner; man or woman practices and private behavioral selections which are related to increased or reduced oral health. (Tolsma D,2009). Dental health behavior has been related to well known fitness behavior, gender, schooling, smoking, alcohol consumption, profits, nutritional habits, physical and social activity, and behavior of parents. (Tada A, Matsukubo T,2003), (Banoczy J et al,2001). Periodontitis consequences from an unbalanced immune reaction of the host in opposition to plaque biofilm. Periodontal pathogenic microorganism secret metabolites and enzymes that worsen tissue harm and, on the same time, leukocytes and fibroblasts produce numerous anti-inflammatory mediators, together with cytokines, prostaglandins, reactive oxidative

species (ROS), protolithic enzymes and metalloproteinase (O'Keefe JH et al,2008). Previous researches have related dental health with tobacco smoking, alcohol consumption, and physical activities. The buildup of plaque and calculus, related to bad oral hygiene, is one of the factors in the improvement of periodontal disease, there has been evidence of a dose-reaction courting between dental hygiene and periodontal health in several disputes. As a result, both good and bad dental hygiene significantly increase the risk of periodontitis. when compared to proper oral hygiene (Lertpimonchai A,etal ,2017).

Wrong brushing techniques, failure to use interdental and abnormal dental visits will lead to plaque accumulation. in a different study, regular brushing was discovered to have a protective effect on oral health, with rare enamel brushing significantly increasing the risk of periodontal inflammation. (Zimmermann H et al,2013) Vegetables and fruits are major sources of nutrients and have antioxidants, which have a effects on periodontal health. Researches detected that a healthy and balanced diet has positive effects on periodontal health. (Simona S et al,2021) Oral health is always associated with popular fitness and numerous studies inside the beyond have clarified an association between socioeconomic factors and oral health. (Jorgensen EB,2000)

Study objectives: is to established relationship between plaque and gingival rates in lifestyle choices. to a person in the city of Erbil.

METHODS AND MATERIALS

The total sample were consisted of 378 individuals their ages were ≥ 18 and whom in search of dental treatment who attended the only four medical centers of the Ministry of Health, which are represented in of the city of Erbil in different regions.

Individuals interviewed by researcher using a researcher-designed questionnaire The interviewer involved the following variables:

Inhalation of smoke related products. (smokers were categorized as smokers (smoked ≥ 10 cigarettes per day) and non-smokers (Riordan M., Arizona, 2008)

Physical activity Substances in this study were defined as whether to exercise (walking, swimming, jogging, cycling) (Qian, J. Cai et al., 2010)

Sport includes an activity normal physical activity or skill performed according to agreed rules and for recreational purposes: for competition, for personal gain, excellence in skill development or a combination thereof (Ibrahim H Mustafa, et al, 2011)

The nutritional status of the patient was indicated based on

servings meat taking per week and dairy milk products. (butter and yogurt), eggs, vegetables and fruits (WordiQ, 2010).

The medical examinations were included plaque index (Silness J & L  e, 1964) gingival index (L  e H & Silness, 1963) examinations of teeth were selected by (Ramfjord SP, 1959) were used Patients with fewer than 20 teeth were excluded from the study.

Statistical analysis: Data management and statistical analysis: Data will be recorded on a specially designed questions, collected and entered in the computer and then analyzed using appropriate data system which is called statistical package for social science (SPSS) (ANOVA and two independent samples t-test) version 22 and the result will be compared between groups with different variables with statistical significance level of < 0.05 .

Procedures:

The correlation between lifestyle and gum health were determined in 378 patients there ages were ≥ 18 and both genders who attended medical centers in Erbil. All participants were sub divided into three categories for (vegetable, fruit, meat and eggs and butter) while for (smoking, exercise and tooth brushing) they were subdivided into two groups. All participants were under go plaque and gingival examination each group hold 21

individual evenly detached. Table (1): shows descriptive statistics for all parameters containing means, standard deviation, standard error, minimum & maximum values. In more detailed comparative (table 2), a t-test for paired samples for the study parameters when consuming vegetables at a day indicated that the differences in PLI and GI were very significant between sub groups.

Table (1): parameters descriptive Statistics in all study categories

	Indices	subgroups	No	Mean	Min	Max	SD	SE
Vegetable /day	PLI	1-3	21	.762	0	3	.764	.159
		3-6	21	.710	0	2	.616	.125
		≥ 6	21	.867	0	2	.707	.146
	GI	1-3	21	.827	0	2	.649	.133
		3-6	21	.942	0	2	.751	.155
		≥ 6	21	.847	0	2	.766	.159
Fruit/day	PLI	1-3	21	.867	0	3	.831	.174
		3-6	21	.587	0	1	.514	.103
		≥ 6	21	.573	0	2	.689	.142
	GI	1-3	21	.823	0	2	.687	.141
		3-6	21	.733	0	2	.677	.139
		≥ 6	21	.673	0	2	.677	.139
Egg&butter /day	PLI	0-1	21	1.155	0	3	.799	.166
		1-2	21	.872	0	2	.660	.135
		2-3	21	.978	0	3	.865	.181
	GI	0-1	21	1.010	0	2	.728	.166
		1-2	21	.885	0	2	.659	.135
		2-3	21	.871	0	2	.865	.125
Meat/week	PLI	0-1	21	1.119	0	3	.779	.158
		1-2	21	.994	0	3	.823	.179
		2-3	21	.849	0	2	.659	.135
	GI	0-1	21	1.14	0	2	.709	.146
		1-2	21	.991	0	2	.713	.147
		2-3	21	.985	0	2	.727	.150
Smoking /day	PLI	0	21	.755	0	2	.625	.127
		≥ 10	21	.623	0	1	.528	.106
	GI	0	21	.783	0	2	.653	.134
		≥ 10	21	.860	0	2	.742	.154
Exercise /day	PLI	0	21	.832	0	3	.791	.164
		1	21	.640	0	2	.596	.121
	GI*	0	21	.918	0	2	.644	.132
		1	21	.639	0	2	.610	.123
Toothbrush /day	PLI	0-1	21	1.52	0	3	.765	.129
		≥ 2	21	.528	0	2	.561	.123
	GI	0-1	21	1.220	0	3	.744	.154
		≥ 2	21	.598	0	2	.649	.133

Table (2): comparative analysis of vegetable taking/day individuals

indices	vegetables/ day			
		mean	S e	P -value
PLI	1-3&3-6	1.762	.159	.000
	1-3 & ≥ 6	1.710	.125	.000
	3-6& ≥ 6	1.867	.146	.000
GI	1-3&3-6	1.827	.133	.000
	1-3 & ≥ 6	1.942	.155	.000
	3-6& ≥ 6	1.847	.159	.000





Figure1:Plaque &gingival indices in vegetable taking/day

Table (3) test of paired samples in fruit consumption per day clarified that the differences in PLI and GI between all subgroups were very significant.

Table (3): comparative analysis of fruit taking/day.

indices	fruit/day			
		mean	S.e	P-value
PLI	1-3&3-6	0.783	.278	.000
	1-3& ≥ 6	1.11	.343	.000
	3-6& ≥ 6	1.830	.103	.000
GI	1-3&3-6	0.732	.250	.000
	1-3& ≥ 6	1.307	.338	.000
	3-6& ≥ 6	1.830	.123	.000




Figur 2: Plaque&gingival indices according to fruit taking /day


While (table 4) revealed that paired sample t-test for each participant were highly significant for both indices in egg and butter taking per day in all subgroups.

Table (4): comparison analysis in egg& butter taking/day.

indices	egg&butter/w week	mean	S.e	P-value
PLI	1&2	.586	.243	.001
	1&3	.944	.309	.000
	2&3	1.585	.226	.000
GI	1&2	.449	.226	.001
	1&3	1.011	.305	.000
	2&3	1.568	.230	.000



Figur 3:Plaque &gingival indicesin egg& butter taking/day.




Figur4: Plaque &gingival indices in relation to meat taking/week

Table (5) showed that paired sample t-test in meat taking per week for PLI and GI were highly significant between all subgroups.

Table (5): comparing study analysis in meat taking/week.

indices	meat/week	mean	S.e	P-value
PLI	1&2	.552	.240	.003
	1&3	1.049	.302	.000
	2&3	1.509	.239	.000
	1&3	.994	.291	.000
	2&3	1.538	.220	.000



Figur5:Plaque&gingival indices in smoker and nonsmoker/day



Table (7). PLI and the GI between groups that perform daily exercise and those that do not, seem to be significant, differences between the two groups not exercise
Table (7) : compares analysis in individual daily exercise or do were also significant.

smoking/day	indices	mean	S.e	P-value
Smoking& nonsmoking	PLI	.274	.128	.149
Smoking& nonsmoking	GI	.364	.135	.041

Table (6). According to this table, the differences in PLI between both groups were not significant, whereas the differences in GI between both groups were significant

Table (6): a comparison analysis in smoker and nonsmoker/day

smoking/day	indices	mean	S.e	P-value
Smoking& nonsmoking	PLI	.274	.128	.149
Smoking& nonsmoking	GI	.364	.135	.041




Table (8): detailed that the matched test showed statistical significant difference in PLI and GI among persons frequently doing their daily brushing or unpredictably

Table (8): compared study parameters among daily tooth brushing /day individuals.

indices	tooth brushing/day			
		mean	S.e	p-value
PLI	Regular& irregular	.801	.211	0.011
GI	Regular &irregular	.711	.201	.001

DISCUSSION

The oral health is associated with popular general health. When oral health is compromised Many systemic illnesses can be manifested in the oral cavity (Akhtar H. Bokhari,2006)

current study begins that, the amount of plaque and gingivitis were more among persons with compounding vegetable consumption in a day, this is may be because of their bad oral hygiene, and the amount of plaque and gingival health were diminishment with expanded daily fruit

consumption this is because of vitamins and antioxidant contains of fruit in which reduce gingival inflammation. (Iwasaki M etal ,2012), (Muniz F.W etal,2015)

Other debates by (Chapple,I.L. C,2010) explain the results of ingesting of vegetable and fruit in disintegrated capsular shape during non-surgical periodontal therapy in 60 non-smokers they have been blundering from slight and moderate periodontitis. They detected small despite the fact that big deduction declines in periodontal depths.

Besides researches by (Vander Velden,2011) concluded that to confer periodontitis and gingivitis, calcium, vitamin diet D and enough Antioxidants retaining nutrients need to be dispatched like fruit, vegetable and berries.

The current debate concluded that mean plaque score and gingival health were more among 0-1 time daily eating butter and egg, this is because of butter and egg calcium content and vitamin D enhance innate immunity towards inflammation. current study is in accordance with (Dietrich, T,2005) study.

Lowest mean plaque score and gingival score were among persons whom eating meat 2-3 times in week. This is because of the iron and O-mega 3(fish canvas) and nutritional vitamins content of meat have antibacterial and anti-oxidant impact, this is in accordance with (Van der Velden U etal,2011)

Former study by ((Nishida, M etal ,2005) determined that for the averting and therapy of gingivitis and periodontitis enough daily input of natural antioxidants, fish (omega-3PUFAs), diet D and calcium are bore. They decided that vitamin D

might again demark the gingival inflammation through its anti-inflammatory impact.

Exercising data from the third National Health and Nutrition Examination Survey (NHANES III) it has been come out that deceased daily consumption of calcium movable in higher periodontal destruction by (Nishida M etal,2005).

The instant study also intimated that plaque accumulating advanced in nonsmoking while progressive mean of gingivitis was in smoker categories persons un healthy life style have effect on periodontal health the improper brushing forms and harming resultants of smoking. (Chapple, I. ,2007), (Vijay K Chava' 2010).

The connection between smoking and an augmented circumstance of periodontitis decided in some examination (Holde, G.E,2017) must no manner longer befall validated. Smokers tend to have a further adverse life- style in all-around. It skill that the relation of tobacco with periodontal condition was associated not alone to miserable oral hygiene but further to miserable typical lifestyle. While study by (Miyuki K etal,2007) whom decide that smoking display the formal effectiveness on periodontitis hazard among life- style components. Smoking might again affect the host- defense system, which can additionally affect periodontal health. The subsisting study about concluded that higher plaque accumulating have been amongst individuality that

would no longer operating exercise daily besides gingival score had been progressive in people who no longer acting exercise. This is due to some degree in practicing greater oxygen will extend to body.

Several studies decided that decreased physical exercise is associated with periodontal disease (Samnienge, P. et al, 2013), (Torrejon Moya, A et al, 2021)

Current study appears that plaque accumulation was high in subjects that not performing their daily tooth brush, additionally gingival inflammation was advanced in individuals their daily tooth brushing.

Current advisement is in accordance with (S. Akhtar Hussain Bokhari, 2006). In some former advisement, ordinary brushing was once located to have defensive holdings on oral health in occasional tooth brushing

significantly compounding the imminence of periodontal inflammation. (Zimmermann H, 2015)

CONCLUSION

In conclusion, the outcomes of the current study materialized that gingivitis is definitely illiberal is more in individuals with unhealthful life style.

REFERENCES

Aiste Zaborskyte, Egle Bendoraitiene: Oral Hygiene Habits and Complaints of Gum Bleeding Among Schoolchildren in Lithuania. *Stomatologija*, 2003; 59(1):31-36.

Akhtar H. Bokhari: Lifestyle Impact on Oral Health. *Department of Oral Health Sciences* 2006, 22(3).

Albander JM. Global risk factors and risk indicators for periodontal diseases. *Periodontol* 2000 2002;29: 177-206.

Banoczy J, Gintner Z, Dombi C. Tobacco use and oral leukoplakia. *J Dent Educ* 2001; 65: 322-6.

Chapple, I. L., Milward, M. R. & Dietrich, T.: The prevalence of inflammatory periodontitis is negatively associated with serum antioxidant concentrations. *Journal of Nutrition* 2007 :137, 657–664.

Chapple, I. L. C., Milward, M. R., Ling-Mountford, N., Weston, P. et al.: Adjunctive effects of a dietary supplement comprising dried whole fruit, vegetable and

berry juice concentrates on clinical outcomes of treatment of periodontitis. *Federation of American Societies for Experimental Biology Journal* 2010: 24, 540.10

Corbet E. Public Health Aspects of Oral Disorders - Periodontal diseases. In: Pine C and Harris R (eds). *Community Oral Health*. Quintessence Publishing Co. Ltd: 2007, 177-189.

Aiste Zaborskyte, Egle Bendoraitiene: Oral Hygiene Habits and Complaints of Gum Bleeding Among Schoolchildren in Lithuania. *Stomatologija*, 2003; 59(1):31-36.

Akhtar H. Bokhari: Lifestyle Impact on Oral Health. *Department of Oral Health Sciences* 2006, 22(3).

Albander JM. Global risk factors and risk indicators for periodontal diseases. *Periodontol* 2000 2002;29: 177-206.

Banoczy J, Gintner Z, Dombi C. Tobacco use and oral leukoplakia. *J Dent Educ* 2001; 65: 322-6.

Chapple, I. L., Milward, M. R. & Dietrich, T.: The prevalence of inflammatory periodontitis is negatively associated with serum antioxidant concentrations. *Journal of Nutrition* 2007 :137, 657–664.

Chapple, I. L. C., Milward, M. R., Ling-Mountford, N., Weston, P. et al.: Adjunctive effects of a dietary supplement comprising dried whole fruit, vegetable and berry juice concentrates on clinical outcomes of treatment of periodontitis. *Federation of American Societies for Experimental Biology Journal* 2010: 24, 540.10

Corbet E. Public Health Aspects of Oral Disorders - Periodontal diseases. In: Pine C and Harris R (eds). *Community Oral Health*. Quintessence Publishing Co. Ltd: 2007, 177-189.

Dietrich, T., Nunn, M., Dawson-Hughes, B. & Bischoff-Ferrari, H. A: Association between serum concentrations of 25-hydroxyvitamin D and gingival inflammation. *American Journal of Clinical Nutrition* 2005; 82, 575–580.

Holde, G.E.; Oscarson, N.; Trovik, T.A.; Tillberg, A.; Jönsson, B. Periodontitis Prevalence and Severity in Adults: A Cross-Sectional Study in Norwegian Circumpolar Communities. *J. Periodontol.* 2017, 88, 1012–1022

Ibrahim H Mustafa. Badya M Najib.Namer
G.Al Tawel: Association between life style
factors and pulmonary tuberculoses in
Erbil.Zanco J.Med.Sci2011;15 (3) : 11-6.

Iwasaki M., Manz M.C., Taylor G.W., Yoshihara
A.,Miyazaki H. Relations of serum ascorbic acid
and α - tocopherol to periodontal disease. *J. Dent.
Res.* 2012;91:167–172.

Jorgensen EB. Effect of socio-economic and
generalhealth status on periodontal conditions
in old age. *J Clin Periodontol* 2000; 27:83

- Lertpimonchai A, Rattanasiri S, Vallibhakara SA, Attia J, Thakkinstian A. The association between oral hygiene and periodontitis: a systematic review and meta-analysis. *Int Dent J*. 2017;67 (6):332–343.
- Löe H, Silness J. Periodontal disease in pregnancy. I. Prevalence and severity. *Acta Odontol Scand*. 1963; 21: 533-551.
- Löe H. Oral hygiene in the prevention of caries and periodontal disease. *Int Dent J* 2000; 50: 129-139.
- Miyuki K, Muneo T, Nobuko N :2007; May 2007: 78(5) ,859-867.
- Martha E. Nunn. Understanding the etiology of periodontitis-an overview of periodontal risk factors. *Perio 2000*. 2003; 32: 1,11-23
- Maurizio S. Tonetti, Henry Greenwell, Kenneth S. Kornman. Staging and grading of periodontitis: Framework and proposal of a new classification and case definition. *J Periodontol*. 2018;89:159– 172.
- Muniz F.W., Nogueira S.B., Mendes F.L.V., Rösing C.K., Moreira M.M.S.M., de Andrade G.M., de Sousa R.C. The impact of antioxidant agents complimentary to periodontal therapy on oxidative stress and periodontal outcomes: A systematic review. *Arch. Oral Biol*. 2015; 60:1203–1214.
- Nishida, M., Grossi, S. G., Dunford, R. G., Ho, A. W., Trevisan, M. & Genco, R. J. (2000) Calcium and the risk for periodontal disease. *Journal of Periodontology*. 2005: 71, 1057–1066.
- O’Keefe JH, Gheewala NM, O’Keefe JO. Dietary strategies for improving post-prandial glucose, lipids, inflammation, and cardiovascular health. *J Am Coll Cardiol*. 2008;51(3):249–255.
- Pihlstrom BL, Michalowicz BS, Johnson NW. Periodontal diseases. *Lancet* 2005; 366: 1809-1820.
- Qian, J.Cai ,M,Gao ,J.,S.,Xu.,L.,Critchley,J.A. Trends in Smoking and quieting in China from 1993 to 2003:National Health Survey data .*Bulletin of the WHO*.2010;88:769-776.
- Ramfjord SP: indices for prevelance and incidence of periodontal disease. *J Periodontal* .1959 ;30:51-59.
- Riordan M. Arizona Health Information. [home page in the internet]. university of Arizona.2007 updated 2008 Sep 4.
- S.Akhtar Hussain Bokhari: Lifestyle Impact on Oral Health;2006 (22) ;3.
- Samnieng, P.; Ueno, M.; Zaitso, T.; Shinada, K.; Wright, F.A.; Kawaguchi, Y. The relationship between seven health practices and oral health status in community-dwelling elderly Thai. *Gerodontology* 2013, 30, 254–261.
- Simona S, Alessandro P, Giuseppe P, Francesco I, Gaetano I: Dietary Factors Affecting the Prevalence and Impact of Periodontal Disease Clinical, Cosmetic and Investigational Dentistry 2021:13 283–292.
- Silness J, Löe H. Periodontal disease in pregnancy. II. Correlation between oral hygiene and periodontal condition. *Acta Odontol Scand*. 1964; 22: 121-135.
- Socransky SS, Haffajee AD. Dental biofilms: difficult therapeutic targets. *Periodontol 2000* 2002;28:12-55.
- Tada A, Matsukubo T. Relationship between oral health behaviors and general health behaviors in a Japanese adult population. *J Public Health Dent*. 2003; 63:250-4.
- Tolsma D. life style available from [http://health.yahoo.net/galecontent/life style](http://health.yahoo.net/galecontent/life_style) cited 2009, may 5.
- Torrejon Moya, A.;Gonzalez-Navarro, B.; Roca-Millan,E.; Estrugo-Devesa, A.; López-López,J. Analysis of Healthy Lifestyle Habits and Oral Health in a Patient Sample at the Dental Hospital of the University of Barcelona. *Int. J. Environ. Res. Public Health* 2021, 18,7488
- Van der Velden U, Kuzmanova D, Chapple ILC. Micronutritional approaches to periodontal therapy. *J Clin Periodontol* 2011; 38 (Suppl. 11): 142–158.
- Vijay K Chava : Effect of lifestyle, education and socioeconomic status on periodontal health: 2010 , 1 (1) : 23-26.
- Wordiq. [home page in the internet]. GUN Documentation License. available from [http://www. Wordiq.com.](http://www.Wordiq.com/) / definition /sports. 2010.
- Zimmermann H, Zimmermann N, Hagenfeld D, Veile A, Kim TS, Becher H. Is frequency of tooth brushing a risk factor for periodontitis? A systematic

review and meta-analysis. *Community Dent Oral Epidemiol.* 2015;43(2):116–127.