

## Y-chromosomal Short Tandem Repeat Variation in Kurd, Assyrian, and Armenian populations in Iraq Kurdistan

Sabriya K. Khalid

Department of Biology, Faculty of Sciences, Zakho University, Dohuk, Kurdistan Region, Iraq

Yousif M. Fattah

Department of Biology, Faculty of Sciences, Zakho University, Dohuk, Kurdistan Region, Iraq, [yousif.fattah@uoz.edu.krd](mailto:yousif.fattah@uoz.edu.krd)

Follow this and additional works at: <https://polytechnic-journal.epu.edu.iq/home>



Part of the Life Sciences Commons

### How to Cite This Article

Khalid, Sabriya K. and Fattah, Yousif M. (2023) "Y-chromosomal Short Tandem Repeat Variation in Kurd, Assyrian, and Armenian populations in Iraq Kurdistan," *Polytechnic Journal*: Vol. 12: Iss. 2, Article 16.

DOI: <https://doi.org/10.25156/ptj.v12n2y2022.pp148-157>

This Original Article is brought to you for free and open access by Polytechnic Journal. It has been accepted for inclusion in Polytechnic Journal by an authorized editor of Polytechnic Journal. For more information, please contact [polytechnic.j@epu.edu.iq](mailto:polytechnic.j@epu.edu.iq).

RESEARCH ARTICLE

# Y-chromosomal Short Tandem Repeat Variation in Kurd, Assyrian, and Armenian populations in Iraq Kurdistan

Sabriya K. Khalid<sup>1\*</sup>, Yousif M. Fattah<sup>2</sup>

<sup>1</sup> Department of Biology, Faculty of Sciences, Zakho University, Dohuk, Kurdistan Region, Iraq

<sup>2</sup> Department of Biology, Faculty of Sciences, Zakho University, Dohuk, Kurdistan Region, Iraq

**\*Corresponding author:**

Yousif M. Fattah,  
Department of Biology,  
Faculty of Sciences,  
Zakho University, Dohuk,  
Kurdistan Region, Iraq.

**E-mail:**

yousif.fattah@uoz.edu.krd

**Received:** 14 September 2022

**Accepted:** 26 October 2022

**Published:** 1 February 2023

**DOI**

10.25156/ptj.v12n2y2022.pp148-157

## ABSTRACT

**Background and objectives:** North central Middle Eastern countries Iran, Iraq, Turkey, and Syria all have persistent Kurdish regions. Over thousands of years, several ethnicities have immigrated, settled, or resided in the region, including Turks, Persians, Arabs, Kurds, Armenians, Assyrians, Chechens, and Azeris. The aim of the present study is to use a set of Y-STR to characterize the genetic structure of the Kurd, Assyrian, and Armenian population and build a phylogenetic tree among them.

**Methods:** Eleven Y-chromosome STRs were evaluated in a total of 90 unrelated males from the Kurds, Armenians, and Assyrians populations in the Kurdistan region of Iraq (DYS19, DYS390, DYS393, DYS426, DYS437, DYS439, DYS447, DYS460, DYS461, DYS481, and DYS576). Using a DNA extraction kit, total DNA was isolated from leukocytes. PCR products were run on 8% polyacrylamide gel with a 50bp ladder as DNA marker to size the bands, and silver staining was used to identify the DNA bands. Power Marker V3.25 software was used to determine variety of genetic parameters, including total allele number, allele frequency, gene diversity, polymorphic information content (PIC), and phylogenetic tree was constructed by MEGA-X software.

**Results:** The total number of alleles identified in the three populations was 380. The sizes of the alleles ranged from 87bp to 275bp. The most diverse loci were DYS447 and DYS576 (GD: 0.949), whereas DYS426 showed the least diversity (GD: 0.896). The Phylogenetic tree divided the populations into two main clusters; The Kurdish and Armenian clades in one cluster and the Assyrian in another cluster. Few of dendrogram leaves from the three examined groups were admixed with each other.

**Conclusions:** This study confirms the high-resolution Y-STR typing's ability to discriminate. It concludes that the genetic distance between Kurd and Armenians is less than the genetic distance between the Kurd and the Assyrians, meaning that the Armenians population are genetically closer to the Kurds population.

**Keywords:** Genetic diversity, population genetics of Iraqi Kurdistan, Y-chromosome STRs

## INTRODUCTION

The Kurds are one of the indigenous peoples of the Mesopotamian plains. The adjacent Kurdish areas of Iran, Iraq, Turkey, and Syria are located in the Mid East's northern central region. Various ethnic groups, including Turks, Persians, Arabs, Kurds, Armenians, Assyrians, Chechens, Azaryan and others have immigrated to, established in, or lived there naturally over the centuries. A significant amount of archaeological evidence suggests that this region is the site of the Neolithic transition (Dogan *et al.*, 2017). Assyrians are an ethnic group indigenous to Assyria, a region located in the Middle East (Oppenheim, 1964). The tribal areas that form the Assyrian homeland are parts of present-day northern Iraq, south eastern Turkey, north western Iran, and north eastern Syria (Nisan, 2002). They are speakers of the Neo-Aramaic branch

isolated population who historically inhabited a region in the Near East bounded by the Mediterranean and Black seas and the Caucasus. They have a complex history including a major geographic displacement during World War I (Haber *et al.*, 2016). Armenian is genetically related to Indo-European languages such as Hittite, Sanskrit, Avestan, Greek, Latin, Gothic, and Slavicas. They have an independent branch of the Indo-European language family. The Armenian and the Kurds historically are living together in one empire or divided between the neighboring empires (Martirosyan, 2014). Numerous DNA-based data are utilized to examine the phylogeography, origins, and demographic history. Numerous Y chromosome polymorphism studies have been conducted for forensic purposes, prenatal testing, and human migrations (Jobling and Tyler-Smith 2000;

Quintana-Murci *et al.*, 2001). With a small amount of mutation and gene conversion, the Y-STR markers are passed from generation to generation without recombination (Rozen *et al.*, 2003; Trombetta *et al.*, 2010). These markers aid to understand the geographical population dynamics and its demographic history (Roewer *et al.*, 2005) and also provide data on the male lineage relationship (Lowery *et al.*, 2013). Due to its ability to distinguish between distinct genetic variants and the highly informative Y chromosome STR haplotypes they formed, Y-STR genotyping has become an important tool in forensic investigations. Non-recombining Y chromosomal markers are more sensitive to founder effects and genetic drift, hence Y-STRs are particularly effective in identifying genetic variations between populations (Heraclides *et al.*, 2017; Li *et al.*, 2016). To illustrate, the Indo-European Kangju and Wusun people were displaced by the male-biased westward expansion of the Xiongnu nomads from the eastern steppe, which resulted an extensive

admixture of east Eurasian lineages (Damgaard *et al.* 2018). Following the formation and blending of many nomadic Turkic states, gene flows between various groups of the former Hunnic empire occurred (Damgaard *et al.*, 2018; Gneccchi Ruscone *et al.*, 2021). The aim of the present study is to use a set of microsatellite markers to characterize the genetic structure of the Kurdish, Assyrian, and Armenian populations in Duhok province. Secondly, is to examine various genetic parameters and build a dendrogram among these three populations in order to determine which primers will be most useful for further population genetic studies.

## SUBJECTS AND METHODS

### 1. Sampling

During the period of 1 December 2021 to 1 March 2022, a total of 90 blood samples were obtained from unrelated males from the Kurds, Armenians and Assyrians populations that reside in Duhok province/Kurdistan Region of Iraq. Informed consent

and with approval of the University of Zakho was obtained for all cases. The genealogical data were also obtained from the donors. The age of participants ranged from 15 to 69 years old.

### 2. DNA Extraction

Genomic DNA was extracted from blood samples using the blood Mini Kit provided by a Chinese company called Dongsheng Biotech. In this research, eleven STR primers were employed as shown in table1.

### 3. PCR amplifications

The following were the PCR cycle parameters: one cycle of initial denaturation at 94°C for 5 min, followed by 36 cycles of denaturation at 94 °C for 1 min. Annealing at 56.5 °C for DYS19, DYS390, DYS460 primers; 60 °C for DYS437, DYS439, DYS481; 61°C for DYS426, DYS447, DYS461 and 59.5 for DYS393, DYS576 for 40 seconds. Extension at 72°C for one minute, and then one cycle of two final extensions at 72 °C for seven minutes.

### 4. Typing

The initial step in running the amplified products was on a 1% agarose gel electrophoresis to identify successful amplifications. Then 8% polyacrylamide gel electrophoresis (PAGE) was used to electrophorese the PCR products. PCR products were run with a 50bp ladder DNA as marker to size the bands. Silver staining was used to identify the DNA bands (Bassam and Gresshoff, 2007).

### 5. Data analysis

Power Marker software version 3.25 was used to compute the allele frequencies and other parameters for each locus. The genetic relationship parameters

calculated according to Reynold (1983) statistics. The similarity matrix was used to construct the dendrogram using the unweighted pair group method arithmetic

Khalid

averages (UPGMA) procedure (Sokal and Michener, 1958). Phylogenetic tree construction was carried out using MEGA-X.

**Table 1: Characteristics of Y-STR markers used in the study.**

| <i>STR markers</i> | <i>Primer Sequences</i>                                              | <i>Repeat motif</i>                        | <i>Annealing Tm. °C</i> | <i>Expected Size(bp)</i> | <i>Reference</i>            |
|--------------------|----------------------------------------------------------------------|--------------------------------------------|-------------------------|--------------------------|-----------------------------|
| DYS19              | F; 5'-CTACTGAGTTTCTGTTATAGT-3'<br>R; 5'-ATGGCCATGTAGTGAGGACA-3'      | (TAGA) <sub>n</sub>                        | 56.5                    | 176 - 212                | Naji. 2020                  |
| DYS390             | F; 5'-TATATTTTACACATTTTGGGCC-3'<br>R; 5'-TGACAGTAAAATGAACACATTGC-3'  | (GATA) <sub>n</sub><br>(GACA) <sub>n</sub> | 56.5                    | 201-242                  | Bai <i>et al.</i> ,2016     |
| DYS393             | F; 5'-GTGGTCTTCTACTTGTGTCAATAC-3'<br>R; 5'-AACTCAAGTCCAAAAAATGAGG-3' | (GATA) <sub>n</sub>                        | 59.5                    | 109-133                  | Butler <i>et al.</i> , 2002 |

|        |                                                                         |                                              |      |           |                      |
|--------|-------------------------------------------------------------------------|----------------------------------------------|------|-----------|----------------------|
| DYS426 | F; 5'-CTCAAAGTATGAAAGCATGACCA-3'<br>R; 5'-GGTGACAAGACGAGACTTTGTG-3'     | (GTT) <sub>n</sub>                           | 61   | 92-98     | Butler et al., 2002  |
| DYS437 | F; 5'-GACTATGGGCGTGAGTGCAT-3'<br>R; 5'-AGACCCTGTCATTACAGATGA-3'         | (TCTA) <sub>n</sub>                          | 60   | 195-229   | Bai et al., 2016     |
| DYS439 | F; 5'-TCGAGTTGTTATGGTTTTAGGTCT-3'<br>R; 5'-GTGGCTTGGAATTCTTTTACCC-3'    | (AGAT) <sub>n</sub>                          | 60   | 198-225   | Butler. 2006         |
| DYS447 | F; 5'-GGTCACAGCATGGCTTGGTT-3'<br>R; 5'-GGGCTTGCTTTGCGTTATCTCT-3'        | (TAATA) <sub>n</sub><br>(TAAAA) <sub>1</sub> | 61   | 206-241   | Redd et al., 2002    |
| DYS460 | F; 5'-CAAATTGCCAACTCTTTC-3'<br>R; 5'-TCTATCCTCTGCCTATCATTATTA-3'        | (ATAG) <sub>n</sub>                          | 56.5 | 162-182   | Bosch et al., 2002   |
| DYS461 | F; 5'-AGGCAGAGGATAGATGATATGGAT-3'<br>R; 5'-TTCAGGTAAATCTGTCCAGTAGTGA-3' | (TAGA)<br>CAGA                               | 61   | 174 - 190 | (NIST) USA in 2017   |
| DYS481 | F; 5'-CTGTTTGAGAGTGTGCGAGA-3'<br>R; ACCCAAGAAGAGCCACACAG-3'             | (CTT) <sub>n</sub>                           | 61   | 119-164   | Moon et al., 2022    |
| DYS576 | F; 5'-TTGGGCTGAGGAGTTCAATC-3'<br>R; 5'-GGCAGTCTCATTTCTGGAG-3'           | (AAAG) <sub>n</sub>                          | 59.5 | 183-207   | Geppert et al., 2009 |

## RESULTS

Using power marker V3.25 software, molecular diversity and population genetic structure such as mean allele number, allele frequency, gene diversity, polymorphic information content, and genetic distance

were analysed with three different populations Kurds (n=30), Armenians (n=30) and Assyrians (n=30). The total number of alleles identified in the three populations was 380 alleles. The sizes of the alleles ranged from 87 to 275bp as shown in Table 2.

**Table 2: Range of allele size of three population Kurd, Armenians and Assyrians**

| Primer |          | Range of allele size, bp | Primer                      |          | Range of allele size, bp |
|--------|----------|--------------------------|-----------------------------|----------|--------------------------|
| DYS19  | Kurd     | 185—219                  | DYS447                      | Kurd     | 208—254                  |
|        | Armenian | 175—197                  |                             | Armenian | 221—275                  |
|        | Assyrian | 176—195                  |                             | Assyrian | 219—268                  |
| DYS390 | Kurd     | 195—240                  | DYS460                      | Kurd     | 165—183                  |
|        | Armenian | 181—200                  |                             | Armenian | 162—190                  |
|        | Assyrian | 202—233                  |                             | Assyrian | 160—178                  |
| DYS393 | Kurd     | 106—127                  | DYS461                      | Kurd     | 163—183                  |
|        | Armenian | 118—133                  |                             | Armenian | 165—180                  |
|        | Assyrian | 118—136                  |                             | Assyrian | 173—190                  |
| DYS426 | Kurd     | 88—100                   | DYS481                      | Kurd     | 200—246                  |
|        | Armenian | 88—98                    |                             | Armenian | 208—250                  |
|        | Assyrian | 87—98                    |                             | Assyrian | 225—273                  |
| DYS437 | Kurd     | 170—186                  | DYS576                      | Kurd     | 175—208                  |
|        | Armenian | 168—200                  |                             | Armenian | 167—204                  |
|        | Assyrian | 153—176                  |                             | Assyrian | 173—192                  |
| DYS439 | Kurd     | 200—225                  | Range of all allele size bp |          | 87 — 275                 |
|        | Armenian | 166—220                  |                             |          |                          |
|        | Assyrian | 191—225                  |                             |          |                          |

The number of alleles per locus in the Kurd population ranged from 7 alleles at locus DYS437 to 16 alleles at locus DYS390, with an average of 10.91 alleles per locus. Allele frequency varied from 0.133 in DYS19 and

DYS576 to 0.333 in DYS437 with mean of 0.197. The gene diversity ranged from 0.798 in DYS437 to 0.922 in DYS576 with a mean of 0.873, indicating a high level of diversity as shown in Table 3.

**Table 3: Allele frequency, Allele number, availability, Gene Diversity, PIC in the Kurd population.**

| <i>Marker</i> | <i>Allele. Frequency</i> | <i>Sample Size</i> | <i>Allele No</i> | <i>Availability</i> | <i>Gene Diversity</i> | <i>PIC</i>  |
|---------------|--------------------------|--------------------|------------------|---------------------|-----------------------|-------------|
| DYS19         | 0.133                    | 30                 | 14               | 1                   | 0.916                 | 0.909       |
| DYS390        | 0.167                    | 30                 | 16               | 1                   | 0.916                 | 0.910       |
| DYS393        | 0.267                    | 30                 | 9                | 1                   | 0.829                 | 0.808       |
| DYS426        | 0.200                    | 30                 | 8                | 1                   | 0.858                 | 0.841       |
| DYS437        | 0.333                    | 30                 | 7                | 1                   | 0.798                 | 0.772       |
| DYS439        | 0.167                    | 30                 | 10               | 1                   | 0.873                 | 0.860       |
| DYS447        | 0.167                    | 30                 | 12               | 1                   | 0.889                 | 0.879       |
| DYS460        | 0.200                    | 30                 | 9                | 1                   | 0.856                 | 0.839       |
| DYS461        | 0.233                    | 30                 | 11               | 1                   | 0.871                 | 0.859       |
| DYS481        | 0.167                    | 30                 | 9                | 1                   | 0.878                 | 0.865       |
| DYS576        | 0.133                    | 30                 | 15               | 1                   | 0.922                 | 0.917       |
| <b>Mean</b>   | <b>0.197</b>             | <b>30</b>          | <b>10.91</b>     | <b>1</b>            | <b>0.873</b>          | <b>0.86</b> |

In Armenian population, the number of alleles per locus ranged from 8 at DYS426 to 18 alleles at DYS447 locus with average of 12.18 alleles per locus. The range of allele frequency ranged from 0.133 in DYS447 and

DYS576 to 0.3 in DYS390 and DYS461 with mean of 0.212. The gene diversity ranged from 0.829 in DYS461 to 0.929 in DYS447 with mean of 0.876 which is higher than Kurds population as shown in Table 4.

**Table 4: Allele frequency, Allele number., availability, Gene Diversity, PIC in the Armenian population**

| <i>Marker</i> | <i>Allele Frequency</i> | <i>Sample Size</i> | <i>Allele No</i> | <i>Availability</i> | <i>Gene Diversity</i> | <i>PIC</i>   |
|---------------|-------------------------|--------------------|------------------|---------------------|-----------------------|--------------|
| DYS19         | 0.233                   | 30                 | 14               | 1                   | 0.869                 | 0.857        |
| DYS390        | 0.300                   | 30                 | 12               | 1                   | 0.853                 | 0.841        |
| DYS393        | 0.200                   | 30                 | 12               | 1                   | 0.882                 | 0.871        |
| DYS426        | 0.233                   | 30                 | 8                | 1                   | 0.851                 | 0.834        |
| DYS437        | 0.233                   | 30                 | 10               | 1                   | 0.853                 | 0.837        |
| DYS439        | 0.167                   | 30                 | 14               | 1                   | 0.898                 | 0.889        |
| DYS447        | 0.133                   | 30                 | 18               | 1                   | 0.929                 | 0.924        |
| DYS460        | 0.167                   | 30                 | 11               | 1                   | 0.889                 | 0.878        |
| 8DYS461       | 0.300                   | 30                 | 9                | 1                   | 0.829                 | 0.809        |
| DYS481        | 0.233                   | 30                 | 11               | 1                   | 0.873                 | 0.861        |
| DYS576        | 0.133                   | 30                 | 15               | 1                   | 0.909                 | 0.902        |
| <b>Mean</b>   | <b>0.212</b>            | <b>30</b>          | <b>12.18</b>     | <b>1</b>            | <b>0.876</b>          | <b>0.864</b> |

In Assyrians population, number of alleles per locus ranged from 8 at DYS426 to 15 alleles at DYS19 and DYS390 locus with average of 11.545 alleles per locus. The allele frequency ranged from 0.133 in DYS19 and

DYS393 to 0.34 in DYS439 with mean of 0.215. The gene diversity ranged from 0.824 in DYS461 to 0.907 in both DYS447 and DYS447 with mean of 0.867 which is lower than Kurds and Armenians population as shown in Table 5.

**Table 5: Allele frequency, Allele No., availability, Gene Diversity, PIC in the Assyrian population**

| Marker      | Allele Frequency | Sample Size | Allele No    | Availability | Gene Diversity | PIC          |
|-------------|------------------|-------------|--------------|--------------|----------------|--------------|
| DYS19       | 0.133            | 30          | 15           | 1            | 0.907          | 0.899        |
| DYS390      | 0.267            | 30          | 15           | 1            | 0.880          | 0.871        |
| DYS393      | 0.133            | 30          | 12           | 1            | 0.893          | 0.884        |
| DYS426      | 0.233            | 30          | 8            | 1            | 0.829          | 0.807        |
| DYS437      | 0.200            | 30          | 11           | 1            | 0.880          | 0.868        |
| DYS439      | 0.300            | 30          | 11           | 1            | 0.831          | 0.813        |
| DYS447      | 0.167            | 30          | 14           | 1            | 0.907          | 0.899        |
| DYS460      | 0.267            | 30          | 9            | 1            | 0.833          | 0.813        |
| DYS461      | 0.267            | 30          | 9            | 1            | 0.824          | 0.803        |
| DYS481      | 0.167            | 30          | 11           | 1            | 0.884          | 0.873        |
| DYS576      | 0.233            | 30          | 12           | 1            | 0.871          | 0.859        |
| <b>Mean</b> | <b>0.215</b>     | <b>30</b>   | <b>11.55</b> | <b>1</b>     | <b>0.867</b>   | <b>0.854</b> |

In order to obtain accurate data analysis, the value of availability (number of observed alleles per number of individuals sampled) was calculated and found to be high, with an average of 1.00 in all the three populations as shown in Tables 3,4, and 5. The polymorphic information content (PIC) values for overall genetic variability also calculated for all primers and in all three populations as shown in Tables 3,4, and 5. The values ranged from 0.887 for the least informative marker, DYS426, to 0.947 for the most informative marker, DYS447. Phylogenetic analysis results are shown in Figure 1. The dendrogram which separated the populations into two main cluster, Assyrian cluster and Kurd-Armenian cluster. The latter, in turn, is divided into two sub-cluster: The Kurd and the Armenians except few individuals were admixed with another cluster or sub-cluster from all the three populations (figure No.1).

## DISCUSSION

In the present study, the Y-STRs were used to determine the allele frequency and genetic variation in 11 loci among Kurds, Armenians and Assyrians populations in Duhok province. The results revealed that within a total of 380 alleles their sizes range 87 to 275 bp as shown in Table 2. These results are similar to those previously reported for the Iraqi Arab families lives in middle Euphrates that their PCR product size of the DYS390

locus was 200 bp, DYS392 locus ranged 93 to 125 bp, DYS393 locus was up to 100 bp and DYS19 ranged from 176 to 212 bp (Naji, 2020). Mean number of alleles per locus scored in this study (Kurd 10.91, Armenian 12.18 and Assyrian 11.55 alleles) showed to be higher than those published in fact sheet of National Institute of Standards and Technology (NIST) USA with an average of 9 alleles per locus. High number of alleles per each population suggests high amount of genetic diversity in the population. Fattah *et al* (2019) reported that the average number of alleles in Kurd population was 5.125, while in this study the number of alleles in Kurd population (10.91) was much higher, this may be due to the use of larger number and different primers. High number of alleles within each population indicates a great level of genetic diversity. The allele frequency in the three groups, Kurd, Armenian and Assyrian was not similar to each other. A study by Ohied and Al Badran (2022) in Basrah population with many similar primers used, showed high allele frequency in all studied loci. Another study by Imad and his colleagues (2013) in middle and south of Iraq population, five of the primers they used were similar to those been used in this study. Allele frequencies in all loci were higher than the results in this study except at DYS439 the value was 0.299. The data in tables 3, 4 and 5 indicate that the mean value of gene diversity in Arminian population is the highest (0.876) then followed by the mean gene diversity in the

Kurd population (0.873) then in Assyrian Khalid  
population



(0.867). Gene diversity in this study was much higher than those reported by Imad and his colleagues (2013) and Naji (2020), while it was similar to those results of Albarzinji (2022) who reported genetic diversity value between 0.848 to 0.392 in Sorani Kurds. In Northern Greece, similar genetic diversity value of 0.9992 also has been scored in 17 Y STR loci, five of these STRs were similar to those used in this study (Leda *et al.*, 2008). The results also revealed that the genetic diversity in Armenian population was higher than those in Kurd and Assyrian populations as shown in Tables 3, 4, and 5. These variations in genetic diversity values in different populations may be attributed to the gene flow and migration during different time of the history. The Polymorphism Information Content (PIC) value is often used to measure the informativeness of a genetic marker. According to Botstein *et al.* (1984), values of PIC greater than 0.5 ( $PIC > 0.5$ ) are considered as highly informative primer. In this study the value ranged from 0.772 at DYS437 locus with 7 alleles in Kurd population to 0.924 at DYS447 locus with 18 alleles in Armenian population. All these primers used in this study therefore can be considered as informative due to their high values. These results are in agree with those of Fattah *et al.* (2019) whom they reported high PIC values. Naji (2020) as well found that DYS19 and DYS392 primers were the most polymorphic in compare to other primers. To evaluate the genetic differentiation and the distance between different populations, a phylogenetic tree was constructed. The phylogenetic tree as shown in figure 1 separated the populations into two major clusters; Assyrian group in one cluster and Kurd-Armenian groups in another cluster. The latter, in turn, is divided into two sub-clusters: The Kurd and the Armenians, except few individuals from one clad clustered to another clad in all the three populations. The genetic distance in three populations indicates that the genetic distance between Kurd population and the Armenians population is less than the genetic distance between the Kurd population and the Assyrians population, meaning that the Armenians are genetically closer to the Kurds. This suggesting long shared history. These groups are sharing the same homeland since thousands of years as well both populations are Indo-European nations while the Assyrian people are Semitic. Admixing of some individuals from one population to another population can be attribute to the long sharing history of living with other for thousands of years. Also, wars, genocides, immigrations and gene flow have its role in admixing some of the individuals from clusters. Another explanation of this, that there will be always an unknown number of males with a similar Y-STR profile (de Knijff, 2022). Tomory *et al.* (2007) in their study demonstrated that there has not been much genetic separation among Hungarian-speaking communities in

the Carpathian Basin. The Hungarian gene pool has been impacted by migration and neighboring gene flow. Therefore, the gene flow may be one of the reasons of the admixture pattern results among these three populations.




Figure 1: Phylogeny relationship of Kurd, Armenian, and Assyrian population using 11 Y STR- markers. The red color are Kurd individuals; the black color is Armenian individuals while the blue color are Assyrian individuals.

## CONCLUSION

All investigated loci have high power of discriminating values, indicating that a DNA-based database can be created using these loci. DYS426 had the lowest gene diversity. DYS447 and DYS576 had a high degree of genetic diversity. The Kurdish community may use these 11 STR loci as a crucial tool for paternal typing and forensic identification based on statistical parameters. The phylogenetic tree separated the populations into two major clusters. Assyrian group in one cluster and Kurd-

Armenian group in another cluster. The latter, in turn, is divided into two sub-cluster: the Kurd and the Armenians, which indicate that the Armenians and the Kurd genetically are close to each other than the Assyrian group.

## REFERENCES

- Albarzinji, B. M. (2022). Population Genetic Study of 17 Y-STR Loci of the Sorani Kurds in the Province of Sulaymaniyah, Iraq. Research square. 1–15. preprint paper.
- Atanasio G, D, E, Marini O, Coppa A, Sellitto D, Trombetta B, Berti A, Cruciani F (2017) Forensic data and microvariant sequence characterization of 27 Y-STR loci analyzed in four Eastern African countries. *Forensic Sci Int Genet* 27:123–131.
- Bai, R., Liu, Y., Li, Z., Jin, H., Tian, Q., Shi, M., & Ma, S. (2016). Developmental Validation of a novel 5 dye Y-STR System comprising the 27 Yfiler Plus loci. *Scientific Reports*, 6, 1–8.
- Bassam, B. J., & Gresshoff, P. M. (2007). Silver staining DNA in polyacrylamide gels. *Nature protocols*, 2(11), 2649.
- Boscha Elena, Andrew C. Leea, Francesc Calafellb, Eduardo Arroyoc, Peter Hennemand, Peter de Knijff d, Mark A. Joblinga, (2002) High resolution Y chromosome typing: 19 STRs amplified in three multiplex reactions. *Forensic Science International* 125, 42–51
- Botstein D, White RL, Skolnick M, Davis RW. Construction of genetic linkage maps in man using restriction fragment length polymorphisms. *Am J Hum Genet*.1984;32:314-31.
- Butler, J. M. (2006). Genetics and genomics of core short tandem repeat loci used in human identity testing. *Journal of Forensic Sciences*, 51(2), 253–265.
- Butler, J. M., Schoske, R., Vallone, P. M., Kline, M. C., Redd, A. J., & Hammer, M. F. (2002). A novel multiplex for simultaneous amplification of 20 Y chromosome STR markers. *Forensic Science International*, 129(1), 10–24.
- Damgaard de Barros P, et al. (2018) The first horse herders and the impact of early Bronze Age steppe expansions into Asia. *Science* 360: eaar7711
- de Knijff, P. (2022). On the Forensic Use of Y-Chromosome Polymorphisms. *Genes*, 13(5), 898.
- Dogan, S., Gurkan, C., Dogan, M., Balkaya, H. E., Tunc, R., Demirdov, D. K., Ameen, N. A., & Marjanovic, D. (2017). A glimpse at the intricate mosaic of ethnicities from Mesopotamia: Paternal lineages of the Northern Iraqi Arabs, Kurds, Syriacs, Turkmens and Yazidis. *PLoS ONE*, 12(11), 1–21.
- Fattah, Y. M., Mohammed, A. B., and Hussien, N. J. (2019). Y-chromosomal STR variation in Kurds and Arabs population in Iraqi Kurdistan. *International Journal of Research in Medical Sciences*, 7(5), 1631.
- Geppert, M., Edelmann, J., and Lessig, R. (2009). The Y-chromosomal STRs DYS481, DYS570, DYS576 and DYS643. *Legal Medicine*, 11(SUPPL. 1), 109–110.
- Gnecchi-Ruscone, G. A., Khussainova, E., Kahbatkyzy, N., Musralina, L., Spyrou, M. A., Bianco, R. A., Radzeviciute, R., Gomes Martins, N. F., Freund, C., Iksan, O., Garshin, A., Zhaniyazov, Z., Bekmanov, B., Kitov, E., Samashev, Z., Beisenov, A., Berezina, N., Berezin, Y., Bíró, A. Z., ... Krause, J. (2021). Ancient genomic time transect from the Central Asian Steppe unravels the history of the Scythians. *Science advances*, 7(13).
- Haber, Marc, Massimo Mezzavilla Yali Xue, David Comas, Paolo Gasparini, Pierre Zalloua and Chris Tyler-Smith (2016). Genetic evidence for an origin of the Armenians from Bronze Age mixing of multiple populations. *European Journal of Human Genetics*. 24, 931–936
- Heraclides A, Bashiardes E, Fernández-Domínguez E, Bertoncini S, Chimonas M, Christofi V, King J, Budowle B, Manoli P, Cariolou MA (2017) Ychromosomal analysis of Greek Cypriots reveals a primarily common pre-Ottoman paternal ancestry with Turkish Cypriots. *PLoS One* 12(6): e0179474.
- Imad H.1 (2013). Genetic variation of 17 Y-chromosomal short tandem repeats (STRs) loci from unrelated individuals in Iraq. *International Journal of Biotechnology and Molecular Biology Research*, 4(8), 119–129.
- Jobling MA, Tyler-Smith C. (2000). New uses for new haplotypes: the human Y chromosome, disease and selection. *Trends Genet* 16(8):356–362.
- Leda K, Jessica L, Jodi A (2008). Population genetics of Y-chromosome STRs in a population of Northern Greeks, *Forensic Sci. Int.* 175(2- 3):250-255.
- Li L, Yu G, Li S, Jin L, Yan S. (2016). Genetic analysis of 17 Y-STR loci from 1019 individuals of six Han populations in East China. *Forensic Sci Int Genet* 20:101–102.
- Lowery RK, Herrera K, Uribe G, Reguiero M, Herrera RJ. (2013). Sub-population structure evident in forensic Y-STR profiles from Armenian geographical groups. *Legal Med* 15(2):85–90.
- Martirosyan, Hrach (2014). Origins and historical

- development of the Armenian language. Report
- Moon, M. H., Hong, S. R., & Shin, K. J. (2022). Sequence variations of 31 y-chromosomal short tandem repeats including 9 rapidly mutating y-chromosomal short tandem repeats analysed by massively parallel sequencing in three u.s. population groups and Korean population. *Journal of Korean Medical Science*, 37(6).
- Naji, Mohammed Zuhair (2020). Study of Some STR Markers of Y Chromosomes in Iraqi families of Middle Euphrates. PhD. Thesis college of science. University of Babylon. Iraq.
- National Institute of Standards and Technology. 100 Bureau Dr, Gaithersburg, MD 20899, USA; 2017.
- Nisan, Mordechai (2002). Minorities in the Middle East: A History of Struggle and Self-Expression, 2nd Edition. McFarland & Co., Jefferson, N.C.
- Ohied, B. M., & Al-Badran, A. I. (2022). Y-chromosome variation in Basrah population. *Journal of Medicine and Life*, 15(2), 202–207.
- Oppenheim A. Leo (1964). *Ancient Mesopotamia* (PDF). University of Chicago Press.
- Quintana-Murci L, Krausz C, McElreavey K (2001) The human Y chromosome: function, evolution and disease. *Foren Sci Inter* 118(2):169–181
- Redd, A. J., Agellon, A. B., Kearney, V. A., Contreras, V. A., Karafet, T., Park, H., De Knijff, P., Butler, J. M., & Hammer, M. F. (2002). Forensic value of 14 novel STRs on the human Y chromosome. *Forensic Science International*, 130(2–3), 97–111.
- Reynolds, J., Weir, B.S., Cockerham, C.C. (1983). Estimation for the coancestry coefficient: basis for a short-term genetic distance. *Genetics* 105: 767-779.
- Roewer L, Croucher PJ, Willuweit S, Lu TT, Kayser M, Lessig R, de Peter K, Jobling MA, Tyler-Smith C, Krawczak M. (2005). Signature of recent historical events in the European Y-chromosomal STR haplotype distribution. *Hum Genet* 116(4):279–291.
- Rozen S, Skaletsky H, Marszalek JD, Minx PJ, Cordum HS, Waterston RH, Richard KW, David C, Page DC. (2003) Abundant gene conversion between arms of palindromes in human and ape Y chromosomes. *Nature*. 423(6942):873–876.
- Sokal, R.R. and Michener, C.D. (1958). *A Statistical Methods for Evaluating Relationships*. University of Kansas Science Bulletin, 38, 1409-1448.
- Tomory, Gyongyver, Bernadett Csanyi, Erika Bogacsi-Szabo, Tibor Kalmar, Agnes Czibula, Aranka Csosz, Katalin Priskin, Balazs Mende, Peter Lango, C. Stephen Downes, and Istvan Rask (2007). Comparison of Maternal Lineage and Biogeographic Analyses of Ancient and Modern Hungarian Populations. *American Journal of Physical Anthropology*, 132(4), 535–544.
- Trombetta B., Cruciani F., Underhill PA., Sellitto D., Scozzari R. (2010). Footprints of X to-Y gene conversion in recent human evolution. *Mol Biol Evol* 27(3):714–725.